

One-loop corrections to the curvature perturbation from inflation

David Seery

Astronomy Unit, School of Mathematical Sciences
Queen Mary, University of London
Mile End Road, London E1 4NS
United Kingdom

E-mail: D.Seery@qmul.ac.uk and djs61@cam.ac.uk

Submitted to: *JCAP*

PACS numbers: 98.80.-k, 98.80.Cq, 11.10.Hi

Abstract. An estimate of the one-loop correction to the power spectrum of the primordial curvature perturbation is given, assuming it is generated during a phase of single-field, slow-roll inflation. The loop correction splits into two parts, which can be calculated separately: a purely quantum-mechanical contribution which is generated from the interference among quantized field modes around the time when they cross the horizon, and a classical contribution which comes from integrating the effect of field modes which have already passed far beyond the horizon. The loop correction contains logarithms which may invalidate the use of naïve perturbation theory for cosmic microwave background (CMB) predictions when the scale associated with the CMB is exponentially different from the scale at which the fundamental theory governing inflation is formulated.

Keywords: Inflation, Cosmological perturbation theory, Physics of the early universe, Quantum field theory in curved spacetime.

1. Introduction

For some years, the theory of inflation [1, 2, 3, 4, 5, 6, 7] has represented our most successful approach to the very early universe. Although we do not know with certainty what physical conditions applied in the very distant past, it is believed that a number of fundamental properties of our universe were determined at that time. Among the most important of these is the ensemble of small density perturbations which later condensed into galaxies and gave rise to the network of cosmic structure we see today. Therefore, one question which must be answered by any theory which purports to describe the very early universe concerns the origin and nature of these perturbations. In inflationary models, the answer is provided by use of an era of quasi-exponential expansion to generate small fluctuations from the vacuum. Remarkably, the properties of these perturbations can be calculated by applying a minimal extension of quantum field theory to the expanding universe [8, 9, 10, 11, 12, 13, 14, 15].

The inflationary perturbations are generated by quantum-mechanical effects among field modes of wavelength k around the time when such modes are of order the Hubble scale, that is, $k \sim aH$. If the inflationary expansion is still ongoing when these perturbations are generated then the local Hubble scale continues to contract, so that eventually they are far outside the horizon, giving $k \ll aH$. In this regime the field behaves in an approximately classical fashion [16]. Therefore the evolution of the curvature perturbation can be calculated reliably until the end of inflation, when it is generally assumed to settle down to a time-independent final value, unless isocurvature modes exist which drive its evolution on large scales [17, 18, 19]. In this simple picture we have a robust prediction of the properties of the perturbations regardless of the details of the inflationary era.

The above account neglects at least one possible source of trouble, namely the question of quantum mechanical corrections. In any quantum theory, including inflation, we are obliged to study the role of so-called *loop corrections*, which may modify or destroy the leading semiclassical prediction. Loop corrections are systematic corrections in powers of a coupling constant, which is small if perturbation theory is to be a good approximation. However, perturbation series are generally asymptotic rather than convergent, and if the coefficients of higher-order terms are large then such corrections need not be negligible even if the coupling constant is small. In such circumstances, these corrections must be taken into account when performing accurate comparisons of theory with observational data; for example, their use is routine when interpreting the outcome of particle physics experiments. Modern cosmological datasets have become remarkably precise [20, 21, 22], and observational improvements imply that future datasets will possess even greater constraining power. It is important that our theoretical predictions keep pace with observational developments. For this reason, as in particle physics, we should ask whether loop corrections can modify the predictions we use to compare with experiment.

If such corrections are to play a role, what form should we expect them to

take? Models of inflation make predictions concerning the properties of the primordial curvature perturbation, ζ . This perturbation is communicated to the cosmic microwave background (CMB) at decoupling. The statistical properties of ζ are predicted by supposing that the universe is geometrically close to a patch of the classical, eternal de Sitter universe, in which the expansion is exactly exponential. The de Sitter universe has a special simplicity as a solution of the Einstein equations since it is maximally symmetric, implying that its curvature is constant. However, as an arena for physics the de Sitter universe is considerably less simple [23]. A horizon surrounds any freely falling observer and screens a portion of the spacetime from view. This horizon is associated with a finite entropy, S_{dS} , which can be thought of as representing a finite state space of size $\sim \exp S_{\text{dS}}$ available to each inertial observer. One therefore expects that perturbative predictions made over a volume $V \gtrsim \exp S_{\text{dS}}$ (in Planck units) are untrustworthy, since the volume V carries more perturbative degrees of freedom than can actually be available in the full quantum gravity [24, 25]. An inflationary epoch lasting for N e-folds causes the universe to expand by a factor e^N , so we expect that the inflationary era cannot be described in perturbation theory for an arbitrarily long time; in order not to violate the volume bound, it is necessary that $N \lesssim S_{\text{dS}}$ for reliable calculations. Related limits on N of various sorts have been reported in the literature [26, 27, 28].

The above intuitive argument implies that we should expect the coefficients in the loop expansion to become large when $N \gg 1$. For sufficiently large N , we will have a breakdown of perturbation theory and thus an apparent loss of predictivity from the early universe. The purpose of the present paper is to explore the details of this loss of predictivity, quantified as a loop correction to the observable perturbation ζ . Although a proper understanding of the breakdown of perturbation theory is important as a matter of principle, it is also important in practice. The apparent success of the inflationary prediction means that it has become routine for experiments which measure temperature anisotropies in the cosmic microwave background to report their findings in terms of constraints on inflationary models. These constraints are usually derived using tree-level predictions for ζ which ignore the possibility of a breakdown in perturbation theory.

The important ramifications which attach to a failure of the perturbative description mean that loop corrections have received attention from many authors, beginning with 't Hooft & Veltman [29]. The possibility of an infrared perturbative breakdown was explored by Sasaki, Suzuki, Yamamoto & Yokoyama [30, 31], and later authors computed loop corrections in a variety of different settings; for example, Refs. [32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53]. We will derive an estimate for the loop correction for an arbitrary potential, using the slow-roll approximation to control the calculation. A central feature of this analysis is the use of the non-linear δN expansion [54, 55, 56, 57]. This expansion provides a systematic method for computing the evolution of ζ (and its correlation functions) in terms of the field perturbations at earlier times (and their correlation functions). Moreover, as will be described below, the δN formulation has many useful properties for the purposes of loop calculations. In

order to apply this method consistently, it is necessary to account for the presence of loop corrections in the field perturbations at early times. The requisite loop correction has recently been calculated [58], and in this paper all these corrections are assembled into a loop correction for ζ , which allows an estimate of the onset of any breakdown in perturbation theory in a number of popular models of inflation.

This paper is organized as follows. In §2 the non-linear δN formula is briefly reviewed, and in §3 it is used to derive the complete one-loop correction to the power spectrum of ζ . Our result agrees with that of Byrnes *et al.*, who recently gave an expression for the loop correction valid to two loops. A feature of the δN expansion is that one is free to construct the initial hypersurface at any time after the relevant $\mathbf{k}$ -modes have left the horizon. This freedom is discussed in §3.3. In §4 we use the formal δN expression to compute an estimate for the loop correction. The loop correction is shown to be divergent on large scales in §4.1, and must be regularized by restricting the computation to some finite box of size ℓ . In §§4.2–4.3 estimates for the regularized loop correction are given in two approximations. In §4.2 the spectrum is assumed to be defined by a constant tilt, whereas in §4.3 we specialise to the case of a monomial potential and compute an explicit, but model-dependent, estimate. Finally, we conclude with a discussion of these results in §5.

We use natural units in which the speed of light c , Planck's constant $\hbar$ and the Planck mass $M_{\text{P}} \equiv (8\pi G)^{-1/2}$ are set equal to unity. The metric signature is $(-, +, +, +)$ and the unperturbed background metric is taken to be of Friedmann–Robertson–Walker form

$$ds^2 = -dt^2 + a(t)^2 d\mathbf{x} \cdot d\mathbf{x}, \quad (1)$$

where t is cosmic time and a is the scale factor of the universe. It is often more convenient to use a conformal time variable, η , defined by $\eta \equiv \int_{-\infty}^t dt/a(t)$. The Hubble parameter satisfies $H \equiv \dot{a}/a$, where an overdot denotes a derivative with respect to t , and inflation occurs whenever $\epsilon \equiv -\dot{H}/H^2 < 1$.

Our theory will consist of Einstein gravity coupled to a scalar field ϕ with potential $V(\phi)$, which can be arbitrary except that it must support an epoch of inflation for some values of the field. The background is taken to be homogeneous with spatially-dependent perturbations $\delta\phi$ which satisfy the smallness condition $|\delta\phi| \ll |\phi|$. When $\epsilon \ll 1$, the field ϕ rolls only a short distance in a Hubble time, and therefore this is known as the slow-roll regime. When slow-roll applies, it is often a good formal approximation to compute in powers of ϵ or related small quantities obtained by taking dimensionless time derivatives of ϵ , although a key theme of this paper will be that such expansions are usually asymptotic rather than convergent. The most important related parameter is η , not to be confused with the conformal time, and defined by $\eta \equiv \ddot{H}/H\dot{\phi}$. This measures the local curvature in the potential V , and in typical models it is small whenever ϵ is, although this need not be true in general.

2. The δN formula

Our theories which govern cosmological evolution at high energies are typically written in terms of a number of microscopic fields whose quanta are the particle species which populate the universe. The interesting cosmological observables are expectation values of products of these fields, often in the special case where the fields which participate in the expectation value are evaluated at equal times but at distinct spatial positions $\{\mathbf{x}_1, \dots, \mathbf{x}_n\}$. It is usually more convenient to translate to Fourier space, and work instead with expectation values of fields evaluated at distinct wavenumbers $\{\mathbf{k}_1, \dots, \mathbf{k}_n\}$.

The power spectrum and bispectrum. The lowest-order expectation value is the one-point function,

$$\langle \delta\phi(\mathbf{k}) \rangle = (2\pi)^3 \delta(\mathbf{k}) O, \quad (2)$$

where O is a dimensionless quantity which depends on the time of evaluation. At tree-level the one-point function is always zero if we have chosen to work in a perturbatively stable vacuum, but it is possible that a non-zero O may be generated radiatively. It was shown in Ref. [58] that to one-loop order O is given by a renormalization-scheme dependent number, which can be absorbed into a redefinition of the background field.

The next-order expectation value is the two-point function,

$$\langle \delta\phi(\mathbf{k}_1) \delta\phi(\mathbf{k}_2) \rangle = (2\pi)^3 \delta(\mathbf{k}_1 + \mathbf{k}_2) P(k_1), \quad (3)$$

where $P(k)$ again depends on the time of evaluation and is known as the power spectrum. It is sometimes more convenient to work in terms of the “dimensionless” power spectrum $\mathcal{P}$, which is related to P by the rule $\mathcal{P} \equiv k^3 P / 2\pi^2$.

The three-point expectation value can be written

$$\langle \delta\phi(\mathbf{k}_1) \delta\phi(\mathbf{k}_2) \delta\phi(\mathbf{k}_3) \rangle = (2\pi)^3 \delta(\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3) B(k_1, k_2, k_3), \quad (4)$$

where B is referred to as the *bispectrum*, since in typical cases it is proportional to a sum of terms quadratic in P [59, 60]. To express B in terms of a dimensionless quantity, it is conventional to introduce the momentum-dependent parameter f_{NL} [60], analogous to $\mathcal{P}$ and occasionally called the “reduced bispectrum,” which is defined to satisfy [61, 57]†

$$B(k_1, k_2, k_3) \equiv -\frac{6}{5} f_{\text{NL}} \left\{ P(k_1)P(k_2) + P(k_2)P(k_3) + P(k_3)P(k_1) \right\}. \quad (5)$$

Note that B (and f_{NL}) can always be written as a function of the three scalars $\{k_1, k_2, k_3\}$, in terms of which it does not depend directly on the relative orientation of the $\{\mathbf{k}_i\}$; alternatively, one can use the momentum conservation condition $\sum_i \mathbf{k}_i = 0$ to rewrite B as a function of only two of the momenta, but in this case it will explicitly depend on their relative orientation, making B (and f_{NL}) always a function of three independent degrees of freedom.

† Sometimes f_{NL} is defined by writing $\zeta = \zeta_g - \frac{3}{5} f_{\text{NL}} \star \zeta_g^2$ (in coordinate space), where ζ is the full curvature perturbation, ζ_g is a gaussian random field, ‘ $\star$ ’ denotes a convolution and f_{NL} is to be thought of as parametrizing a possible χ^2 -type contribution to ζ . To linear order in f_{NL} , this expansion for ζ in terms of ζ_g produces the formula (5). However, since there is no motivation for such a specific type of non-linearity in ζ , we follow Lyth & Rodríguez [57] in adopting Eq. (5) for f_{NL} , whatever its source.

Tilt. The dimensionless quantities $\mathcal{P}$ and f_{NL} are generally scale-dependent. Their scale dependence can locally be characterized by two *tilt* parameters n and n_{NL} [62], defined by

$$n - 1 \equiv \frac{d \ln \mathcal{P}}{d \ln k} \quad \text{and} \quad n_{\text{NL}} \equiv \frac{d \ln f_{\text{NL}}}{d \ln k}, \quad (6)$$

where in the definition of n_{NL} we take f_{NL} to be evaluated at equilateral momenta $k_i = k$, and by convention $n - 1$ is chosen as the tilt of $\mathcal{P}$ rather than n .[‡] These numbers are typically small [63]. For a limited range of wavenumbers it may be an acceptable approximation to take $\mathcal{P}$ and f_{NL} as almost flat over the range, whereas if it is necessary to characterize $\mathcal{P}$ or f_{NL} over a larger range then one can write

$$\mathcal{P} \approx \mathcal{P}|_{k_0} \left(\frac{k}{k_0} \right)^{(n-1)|_{k_0}} \quad \text{and} \quad f_{\text{NL}} \approx f_{\text{NL}}|_{k_0} \left(\frac{k}{k_0} \right)^{n_{\text{NL}}|_{k_0}} \quad (7)$$

where k_0 is some pivot wavenumber which is characteristic of the range of wavenumbers in question, and ‘ $|_{k_0}$ ’ denotes evaluation at this point. However, both these approximations break down over an exponentially large range of wavenumbers since in general none of $\mathcal{P}$, f_{NL} , n or n_{NL} are constant. Over such an exponentially large range Eq. (7) substantially overpredicts the spectrum, which typically grows only like a power of $\ln k$ rather than a power of k [44, 64]. Where possible we will use Eqs. (6)–(7) to give model-independent estimates for tilted spectra, but it is necessary to make more detailed model-dependent estimates where an exponentially large range of scales applies (see §4.3).

The δN formula. The microscopic fields are not observable by themselves; only a specific combination of them can be observed as the gauge-invariant perturbation ζ [65] which is communicated to the CMB. An extremely powerful and computationally convenient way to obtain this combination is to employ the so-called δN formula [54, 55, 66, 56, 57]. Consider the Arnowitt–Deser–Misner [67] line element

$$ds^2 = -N dt^2 + h_{ij}(dx^i + N^i dt)(dx^j + N^j dt), \quad (8)$$

in which the perturbations are parametrized by a lapse function N , the shift vector N^i and the spatial metric h_{ij} . For any choice of t , we can write the spatial metric on the three-dimensional slices of constant time in the form $h_{ij} = a(t)^2 \exp\{2\psi(t, \mathbf{x})\} \delta_{ij}$.[§] The quantity ψ is known as the *curvature perturbation* associated with this slicing.

[‡] Note that n_{NL} clearly does not capture the most general variation in f_{NL} under a change of the momenta, since f_{NL} is a function of three degrees of freedom. More generally, one could define the tilt of f_{NL} as a vector quantity obtained by taking the gradient with respect to the k_i .

[§] This is not strictly true, because this is not the most general perturbation of a spatial slice. The curvature perturbation ψ amounts to a fluctuation in the trace of h_{ij} , which describes the volume expansion in a local region of the slice. In principle there are further vector and tensor modes which describe vorticity and gravitational waves. These modes are set to zero. In general this is *not* a consistent procedure, because scalar, vector and tensor perturbations mix beyond linear order. Therefore the presence of scalar perturbations will induce perturbations in the vector and tensor modes. However, we expect that these effects will not destroy the leading order prediction for ζ , which we can obtain by focusing simply on ψ .

Consider any two hypersurfaces at times t_* (the *initial* hypersurface) and t_c (the *final* hypersurface), where $t_c > t_*$. In the perturbed universe the number of e-foldings of inflation between these slices will vary from place to place, and satisfies

$$N(t_* \rightarrow t_c, \mathbf{x}) = N_0(t_* \rightarrow t_c) + \psi_c(\mathbf{x}) - \psi_*(\mathbf{x}), \quad (9)$$

where N_0 represents the number of e-folds which would have taken place in the unperturbed background. By choosing the initial slice to be flat, so that $\psi_* = 0$, and the final slice to be uniform density, so that $\psi_c = \zeta_c$, one obtains the simple expression $\zeta = \delta N$, where $\delta N \equiv N(\mathbf{x}) - N_0$ is the shift in e-folds between t_* and t_c generated by the perturbations. We expect widely separated Hubble volumes to evolve locally like independent universes, so on large scales this shift can be computed merely by accounting for the variation in initial conditions from place to place at time t_* ,

$$\zeta_c(\mathbf{x}) = \sum_{n=1}^{\infty} \{\delta\phi_*(\mathbf{x})\}^n \left(\frac{\partial}{\partial\phi_*} \right)^n N(t_c, t_*), \quad (10)$$

where the $\delta\phi_*$ are the field perturbations at t_* , evaluated on the flat slicing. (Note that ζ_c defined by this expression has no dependence on t_* .) In principle N depends on all the initial conditions on the initial slice, and because ϕ is governed by a second-order differential equation it will generally be necessary to specify the pair $\{\phi_*, \dot{\phi}_*\}$ at t_* . We will make the simplifying assumption that the slow-roll approximation applies at the initial time. This relates $\dot{\phi}_*$ to ϕ_* , and hence it is unnecessary to include $\dot{\phi}_*$ separately.

Expectation values. Eq. (10) can be used to compute n -point correlation functions of the curvature perturbation ζ , by first constructing a product of n copies of (10) and taking expectation values $\langle \dots \rangle$ in the appropriate vacuum state. After application of this process, and working to lowest non-trivial order in $\delta\phi$, it follows that

$$\langle \zeta(\mathbf{x}_1) \zeta(\mathbf{x}_2) \rangle_c = \left\{ \frac{\partial N(t_c, t_*)}{\partial \phi_*} \right\}^2 \langle \delta\phi(\mathbf{x}_1) \delta\phi(\mathbf{x}_2) \rangle_*, \quad (11)$$

where the subscript c denotes evaluation on the final slice at time t_c . Eq. (11) shows that we can compute the power spectrum at any late time t_c using our knowledge of correlators only at some fixed early time t_* . By repeating the above process for higher n -point expectation values and to higher orders in $\delta\phi$, it is obvious that we can use the δN formula to generate predictions for any n -point function of ζ to any required accuracy compatible with the neglect of spatial gradients.

The manipulations described above are now standard in the literature. However, the foregoing discussion makes it completely clear that the δN formula is merely an identity which expresses ζ in terms of geometrical quantities. From this point of view, ζ is nothing more than a book-keeping object which describes how much one region of the universe has expanded (if $\zeta > 0$) or contracted (if $\zeta < 0$) relative to the mean expansion. Therefore, as has been observed before, it is a pure gauge mode in exact de Sitter space [61].

How are we to assign an interpretation to the notion of mean expansion? This is necessary to determine what question we actually answer when we compute properties

of ζ using the δN formula. It corresponds to specifying which region of the universe is supposed to be described by the unperturbed background metric (1). In practice, we do not imagine that the inflationary patch which gave rise to the large flat region we observe today fills our entire spatial slice. For example, this may happen because the universe emerged from some primeval phase with initial conditions which were chaotically varying, and the requirements for successful inflation were only realised at a fraction of spatial positions. Arguments of this sort are invoked for chaotic inflationary models, and arise in a similar way for cosmologies based on the so-called “landscape” of string theory vacua. In such scenarios we should expect spacetime to retain a disordered, inhomogeneous and anisotropic character beyond the boundary of the inflationary patch.

It is quite clear that we should only associate the FRW background metric (1) with the region of spacetime containing the primordial inflationary patch, perhaps after a few e-foldings of inflation so that the patch is sufficiently close to spatial flatness and isotropy [68]. One then expects that whatever primordial unpleasantness lurks beyond the boundary is irrelevant when we make predictions from inflation, provided we live sufficiently far from the boundary. This is analogous to the decoupling theorem in field theory (see, *eg.*, [69]). It follows from the simple point of principle, related to cluster decomposition, that in order to make a prediction at one point we should not need to know the disposition of the universe at some far distant location.

In practice this means that $a(t)$ should be interpreted as providing only an effective description within some box which coincides with the inflationary patch. But we are quite free, if we wish, to imagine that we are working within some smaller box, as long as it is large enough to contain scales of order the present horizon size. Whichever box size we pick, ζ represents the modulation in local expansion as one varies one’s location within the box. Do the answers to calculations depend on the box size? Certainly. In fact, $a(t)$ will generally shift in value between differently sized boxes [70, 71, 72, 73], because the split between background and perturbations is different on disparate scales. One can obtain the correct variation in $a(t)$ by demanding that the expectation value of ζ vanish within the box, that is, $\mathbb{E}_\ell(\zeta) = 0$, where $\mathbb{E}_\ell$ represents the spatial expectation value within a box of characteristic size ℓ . This process, which determines the physical perturbation mode within the box, is analogous to imposing a renormalization condition which determines a physical property of a particle, such as its mass.

This ℓ -dependence of $a(t)$ and ζ imply that one is left with predictions for the properties of ζ which depend on the box size, ℓ . As emphasized by Boubekur & Lyth, one can aggregate predictions in boxes of size ℓ into larger boxes of size $\ell' \gg \ell$ in such a way that the dependence of ζ on ℓ is cancelled by the variation of background quantities [71, 73]. This is analogous to renormalization group invariance in field theory. Nevertheless, the statistical properties of ζ itself depend on ℓ . At first sight which seems like a failure of locality. The resolution is that if the properties of ζ calculated within a small box are accessible to experiment, the properties of ζ calculated some much larger box are not, because they require measurements to be made on scales which are larger than our present horizon. In order to make predictions for the CMB, we wish ζ to

represent the perturbation which is actually present in the radiation when the universe becomes transparent. This entails making an optimal choice for ℓ in the same way that making a prediction for a scattering experiment at some characteristic momentum transfer entails making an optimal choice for the running coupling constant.

In what follows we compute the loop correction in $\langle \zeta(\mathbf{k}_1)\zeta(\mathbf{k}_2) \rangle$. This will enable us to discuss the ℓ -dependence of the prediction, and select an appropriate box size for the purposes of CMB predictions.

3. The one-loop correction to the power spectrum of ζ

3.1. A renormalized δN formula

In this section, we compute the formal δN expression for the one-loop correction to $\langle \zeta(\mathbf{k}_1)\zeta(\mathbf{k}_2) \rangle$. This expression will turn out to be divergent and will require, as outlined in §2 above. In order to organize the calculation, it is convenient to express the various contributions to ζ using a set of graphical rules, analogous to the Feynman diagrams of quantum field theory. Similar diagrams were introduced by Crocce & Scoccimarro [74] and adapted for use in the present context by Zaballa, Rodríguez & Lyth [75]. The use of such diagrams has recently been formalized to all orders by Byrnes *et al.* [76] using a slightly different notation. Their principal use is to keep track of various loop corrections which arise from the δN algorithm described above; when Eq. (10) is translated to Fourier space, the products $\delta\phi(\mathbf{x})^n$ at a point $\mathbf{x}$ become convolutions with independent integrals over $n - 1$ momenta. Some of these momenta are removed after taking expectation values, but this cancellation is not always exact and any unconstrained integrals can be thought of as making loop corrections to the expectation value.

The rules for drawing diagrams can be summarised as follows. In order to evaluate the contribution to an n -point correlator, one draws all diagrams with n external legs which inject momenta $\{\mathbf{k}_i\}$ into the diagram, where $i = 1, \dots, n$. To each external leg one attaches a single m -valent vertex (where $m \geq 2$), at which momentum conservation applies. Each m -valent vertex contributes a “coupling constant” $\partial^{m-1}N/\partial\phi^{m-1}$. The internal lines attached to such vertices terminate at any number of special vertices, which we label with a cross ‘ $\times$ ’. One then draws all combinations of j -gons which can be used to connect the $\times$ -vertices with no $\times$ -vertex left over, replacing each j -gon by a j -point expectation value of the $\delta\phi$ as shown in Fig. 1. These expectation values should be evaluated at the momenta which flow into the corresponding $\times$ vertices. For clarity, we allow a special notation for 2-gons. (See Fig. 2.) Finally one integrates over all internal momenta in order to obtain the expectation value. In such diagrams, it is sometimes useful to imagine the various j -gons as representing initial conditions which flow from the interior of the graph to the outside, encountering time evolution at each m -valent vertex.

The discussion in §2 implies that we should generalize the formal δN expansion,




Figure 1. Diagrammatic representation of the δN expansion. To represent an n -point correlator one draws all m -valent vertices attached to n external legs. Internal lines terminate at special vertices marked by a cross ‘ $\times$ ’. One then draws all possible j -gons which connect the $\times$ -vertices, of which two possible examples are displayed. 2-gons are treated specially for notational convenience (see Fig. 2). Each m -valent vertex contributes a “coupling constant” $\partial^{m-1}N/\partial\phi_*^{m-1}$.




Figure 2. Special representation for 2-gon diagrams. In the left-hand diagram the two $\times$ -vertices should be connected by a 2-gon, but this is difficult to draw. Instead, we adopt the notation shown in the right-hand diagram, where the 2-gon is represented by two parallel lines transverse to the propagators in which they are inserted. In principle there should also be 1-gon initial conditions, where a single $\times$ -vertex is tied off at a one-point function $\langle\delta\phi\rangle$, but because we are neglecting the one-point function on large scales such 1-gons do not appear in these diagrams.

Eq. (10), by inserting a zero-momentum expectation value δN_0

$$\zeta = \delta N_0 + \sum_{n=1}^{\infty} \left\{ \delta\phi_*(\mathbf{x}) \right\}^n \left(\frac{\partial}{\partial\phi_*} \right)^n N(t_c, t_*), \quad (12)$$

which is to be adjusted so that $\mathbb{E}_\ell(\zeta) = 0$, meaning that ζ accurately represents the perturbation in expansion over a box of characteristic size ℓ . Enforcing this condition shows that δN_0 must obey

$$\delta N_0 = - \sum_{j=2}^{\infty} \frac{1}{j!} \frac{\partial^j N}{\partial\phi^j} \langle\delta\phi^j\rangle, \quad (13)$$

where we have assumed that $\langle\delta\phi(\mathbf{x})\rangle = 0$ and that statistical homogeneity forces $\langle\delta\phi^j\rangle \equiv \langle\delta\phi^j(\mathbf{x})\rangle$ to be independent of $\mathbf{x}$ for $j \geq 2$. This guarantees that δN_0 carries zero momentum. Angle brackets $\langle\cdots\rangle$ denote an expectation value in the quantum vacuum at past infinity, as usual, and $\mathbb{E}_\ell$ denotes a spatial average over a comoving spherical box of characteristic size ℓ .

The scale factor is not the only background quantity which will receive a renormalization. In general we can expect all parameters which specify the homogeneous background model to each receive renormalizations.[†] However, the presence of δN_0

[†] This is similar to the proposal of Kolb *et al.* [77], who suggested that a related renormalization of the expansion rate *at the present day* could be responsible for the presently observed accelerated expansion. This proposal was criticized on various grounds in Refs. [78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88] and

does not affect n -point expectation values of ζ for $n > 1$, because it leads to disconnected contributions. These are not the contributions we wish to measure from CMB observations and therefore can be discarded in the present context.

3.2. The power spectrum of the primordial curvature perturbation

The simplest possible graph corresponds to Eq. (11), which arises from the product of the two linear factors in (10),

$$\text{---}\text{---}\text{---} \mapsto \left(\frac{\partial N}{\partial \phi_*} \right)^2 \langle \delta\phi(\mathbf{k}_1) \delta\phi(\mathbf{k}_2) \rangle_*, \quad (14)$$

where the subscript ‘*’ denotes evaluation on the initial slice. Although Eq. (14) is clearly a tree diagram in terms of the δN diagrammatic rules described in §3.1, it implicitly contains loop corrections which are present in the initial correlator $\langle \delta\phi(\mathbf{k}_1) \delta\phi(\mathbf{k}_2) \rangle$. This illustrates a general feature which arises in calculating ζ correlators using δN ; there are two distinct types of loop. The first sort are the quantum-mechanical loops (called q-loops in Ref. [16]) present in the expectation values of the $\delta\phi$. The second sort are loops which arise from expanding ζ correlators using the δN algorithm (called c-loops in Ref. [16]). To arrive at a consistent result at any given order in the loop expansion, it is necessary to retain loop contributions from *both* these sources. Therefore, we require the q-loop correction in $\langle \delta\phi(\mathbf{k}_1) \delta\phi(\mathbf{k}_2) \rangle_*$. This q-loop has been computed by Sloth [51, 52] in the limit where t_* is long after the mode with wavenumber $k \sim k_1 = k_2$ exited the horizon, and has recently been given in Ref. [58] in the opposite case where t_* is the horizon-crossing time associated with k . We are free to base the δN expansion on either of these slices, and a discussion of their merits is deferred to §3.3 below.

At whatever time it is evaluated, the q-loop associated with Eq. (14) is accompanied by a number of c-loops arising from δN integrations. The first of these can be written

$$\text{---}\bigcirc\text{---} \mapsto (2\pi)^3 \delta(\mathbf{k}_1 + \mathbf{k}_2) \frac{\partial N}{\partial \phi_*} \frac{\partial^2 N}{\partial \phi_*^2} \int \frac{d^3 q}{(2\pi)^3} B_*(\mathbf{k}_1, \mathbf{k}_2 - \mathbf{q}, \mathbf{q}), \quad (15)$$

where B is the bispectrum of the field fluctuations, defined in Eq. (4), and the subscript ‘*’ again denotes evaluation on the initial slice. This term gives a contribution to the loop correction which depends on the presence of non-gaussianity in the field perturbations $\delta\phi$. We shall see in §4.2 below that this term can in fact give the largest contribution to the loop correction if f_{NL} grows on very large scales, although this does not happen in the usual scenario. The presence of a single unconstrained momentum integral, here labelled $\mathbf{q}$, indicates that this is a one-loop contribution. The finite time of evaluation,

elsewhere. The present discussion differs from that given by Kolb *et al.* because there is no suggestion here that renormalization of the acceleration rate would have experimental consequences for observers within any given horizon volume; indeed, precisely the *opposite* is the case. The renormalization is only relevant for so-called “meta-observers” who can simultaneously compare rates between many different Hubble patches; it is relevant for what we can compute, but not necessarily for what we can observe.

at conformal time η_* , provides a natural upper cut-off in the momentum integral. On scales characterized by wavenumbers *larger* than $q \sim -\eta_*^{-1}$ no perturbations have yet been generated; on these scales, the spectrum and all higher correlation functions such as B and the trispectrum T are zero.[‡] The upper limit on the integral therefore depends on when we take the initial slice to be constructed, whereas the comoving box size ℓ provides a natural (and time-independent) lower cutoff at $q \sim \ell^{-1}$.

The next contribution arises from the product of two tree-level spectra. There are two ways such a product can arise. The first corresponds to the diagram

$$\text{---} \bigcirc \text{---} \mapsto (2\pi)^3 \delta(\mathbf{k}_1 + \mathbf{k}_2) \frac{1}{2} \left(\frac{\partial^2 N}{\partial \phi_*^2} \right) \int \frac{d^3 q}{(2\pi)^3} P_*(|\mathbf{k}_1 - \mathbf{q}|) P_*(r), \quad (16)$$

whereas the second corresponds to a different configuration

$$\text{---} \bigcap \text{---} \mapsto (2\pi)^3 \delta(\mathbf{k}_1 + \mathbf{k}_2) \frac{\partial N}{\partial \phi_*} \frac{\partial^3 N}{\partial \phi_*^3} \int \frac{d^3 q}{(2\pi)^3} P_*(k) P_*(q), \quad (17)$$

and the remarks below Eq. (15) concerning the presence of an upper- and lower-cutoff in the momentum integral apply equally to both these loops.

These diagrams exhaust the possible one-loop contributions if we assume that the one-point function of the field fluctuations is zero, $\langle \delta\phi(\mathbf{k}) \rangle = 0$. As discussed in §2, the one-point function O is given to one-loop order by a renormalization-scheme dependent number which can be absorbed into the background field $\phi(t)$. This number is accompanied by other unknown renormalization-scheme dependent numbers which are present in the two- and higher n -point correlation functions. We will assume that to a good approximation, all such unknown quantities can be ignored; since we will not keep track of scheme-dependent numbers in the higher n -point functions there is no point keeping track of O or a renormalization of the background field $\phi(t)$ either. Therefore, all contributions from one-point insertions will be discounted in the remainder of this paper.

These expressions agree with those recently given by Byrnes *et al.* in Ref. [76].

3.3. Where should the initial slice be set?

In the discussion of §3.2 the initial time at which the δN expansion is to be constructed, η_* , was left arbitrary. The initial time plays two roles in the analysis: it determines whether the field correlators $\langle \delta\phi(\mathbf{k}_1) \cdots \delta\phi(\mathbf{k}_n) \rangle_*$ are to be evaluated at the approximate time when the mode with wavenumber $k \sim k_1 = k_2$ crosses the horizon, or if they are to be evaluated at a much later time; and it provides an upper cut-off in integrals over c-loop momenta. These two effects must combine in such a way that the curvature perturbation is independent of η_* .

[‡] If we were computing these objects using the underlying quantum field theory of ζ rather than by following the δN method, then these wavenumbers would lie in a region of η -integration where the integrand is strongly suppressed by decaying exponentials, leaving almost no net contribution.

The loop correction evaluated long after horizon crossing has been computed by Sloth [51, 52] in terms of the variance $\langle \delta\phi^2 \rangle \equiv \int d\ln q \mathcal{P}(q)$, which is in principle a infrared divergent quantity and should be cut off at a momentum scale of order the box size, $q \sim \ell$, if $\mathcal{P}$ does not vanish as $q \rightarrow 0$. The loop correction is found to diverge essentially logarithmically with the time of observation,

$$\langle \delta\phi(\mathbf{k}_1)\delta\phi(\mathbf{k}_2) \rangle_* \sim (2\pi)^3 \delta(\mathbf{k}_1 + \mathbf{k}_2) P_*(k) \left\{ 1 + \alpha \text{Ci}(-2k\eta_*) \langle \delta\phi^2 \rangle \right\} \quad (18)$$

where $|k\eta_*| \ll 1$ by assumption, α is a numerical factor of order $\mathcal{O}(\epsilon)$, P_* is the tree-level power spectrum, which is assumed to be almost constant between horizon crossing and the time of observation, and $\text{Ci}(x)$ is the real part of the imaginary exponential integral,

$$\text{Ci}(x) \equiv \text{Re} \left\{ \text{Ei}_1(ix) \right\}, \quad \text{where} \quad \text{Ei}_n(z) \equiv \int_1^\infty \frac{dt}{t^n} e^{-tz} \sim -\gamma - i\pi - \ln z \quad \text{for } z \ll 1, \quad (19)$$

where γ is Euler's constant. Eq. (19) apparently contains *two* distinct divergences.

One divergence is implicit in $\langle \delta\phi^2 \rangle$. In exact de Sitter space $\mathcal{P}(q)$ is a constant, so $\langle \delta\phi^2 \rangle$ appears to contain a logarithmic divergence at zero momentum. This is caused a pile-up of modes outside the horizon, arising from gravitational redshifting, and is ultimately traceable to the quasi-de Sitter expansion. As modes are redshifted outside the horizon they carry perturbations, and the ever-increasing number of modes in this infrared phase space leads to large effects. However, the logarithmic divergence itself is fictional because the original inflationary patch was presumably not infinite in extent. Working in a finite box cuts off the divergence on a momentum scale $q \sim \ell^{-1}$.

The second divergence is present in the $\eta_* \rightarrow 0$ limit, and becomes manifest when one observes the correlator long after the $\mathbf{k}$ -mode in question has passed outside the horizon. This divergence comes from the logarithmic term in $\text{Ci}(-2k\eta_*)$. As with the pile-up of modes, this divergence is fictional in the sense that no observer within the universe can really set $\eta_* = 0$. Instead, observations must be made at a finite time and this finite time gives a cut-off on η_* .

Although we do not encounter a genuine infinity in either case, these “divergences” can generate large coefficients in the perturbation series. Consider first the divergence as $\eta_* \rightarrow 0$, which apparently signals a breakdown of perturbation theory at some time of order $N \sim 1/\epsilon$ e-folds after horizon crossing. (Related breakdowns have been reported elsewhere in the literature; see, for example, Refs. [46, 26, 27].) The presence of this term endows the correlator with a time dependence that is not visible at tree-level. Therefore one should worry whether this time dependence could spoil conservation of ζ on superhorizon scales. If so this would be cause for serious concern, because ζ -conservation is known to hold to all orders of the δN formula [56] and therefore cannot be spoiled by c-loops, and since no field correlator is ever evaluated at the time ζ is observed it cannot be spoiled by q-loops among the $\{\delta\phi\}$ either.

In fact, the time dependence in Eq. (19) is merely what is *required* to maintain conservation of ζ . Consider any two widely separated initial times. When evaluated at the same time of observation, the coefficient $\partial N / \partial \phi_*$ evolves between these two initial slices because the classical field ϕ is rolling down its potential. The correlator

$\langle \delta\phi(\mathbf{k}_1)\delta\phi(\mathbf{k}_2) \rangle$ must likewise evolve between the two initial slices to compensate, in order that ζ remains constant. It is this evolution which is represented by the logarithmic “divergence” in (19). Indeed, Eq. (19) should be thought of as the beginning of a Taylor series in $\ln(-k\eta_*)$, with the higher-order terms in the series generated by higher-order loop corrections. This series is of the form studied by Gong & Stewart [89, 90]. In particular, it implies that perturbation theory in powers of slow-roll parameters breaks down at sufficiently late times, of order $N \sim 1/\epsilon$ e-folds after horizon crossing.

Despite this breakdown in perturbation theory, our ability to make predictions from theories of the early universe is not really impaired. In such situations, one would ordinarily attempt to resum the large logarithms using a form of the renormalization group equation. This resummation procedure effectively moves the large coefficients from higher-order terms in the perturbation series into the leading-order term. In the context of Eq. (19), resummation of the large logarithms clearly yields the superhorizon evolution of the correlator. Since the perturbations behave classically in the superhorizon limit, we should expect that this evolution will correspond to the classical evolution of the field [54, 91, 92, 93].

Now consider the alternative choice, where the loop correction associated with a mode of wavenumber $\mathbf{k}$ is evaluated at the time when $\mathbf{k}$ crosses the horizon. At horizon crossing it follows that $-k\eta_* \simeq 1$. The large logarithmic term $\text{Ci}(-2k\eta_*)$, and any higher-order logarithms, are therefore completely negligible. This version of the loop correction has recently been computed in Ref. [58], with the result

$$\langle \delta\phi(\mathbf{k}_1)\delta\phi(\mathbf{k}_2) \rangle_* = (2\pi)^3 \delta(\mathbf{k}_1 + \mathbf{k}_2) P_*(k) \left(1 - \frac{4}{3} \mathcal{P}_* \ln k + \dots \right). \quad (20)$$

The scale which accompanies k inside the logarithm is set by the details of physics in the ultraviolet.[§] Although such corrections are extremely interesting, they are not the focus of the present paper. For this reason we will ignore them in what follows. At this order, $\langle \delta\phi(\mathbf{k}_1)\delta\phi(\mathbf{k}_2) \rangle_*$ is free of infrared divergences.

The key lesson to draw from a comparison of Eqs. (19) and (20) is that the divergence at late times, associated with powers of $\ln(-k\eta_*)$, has been resummed into the δN coefficient associated with this correlation function.^{||} It is a remarkable virtue of the δN formulation that this resummation is implicit in our freedom to fix the initial slice at any convenient time, without any necessity to deploy a complex formalism based on the renormalization group. Eqs. (19) and (20) are therefore merely different representations

[§] In v1–v2 of the `arXiv` version of this paper, and the version which subsequently appeared in JCAP, the regulating scale which accompanies the logarithm was erroneously identified as the infrared scale ℓ . I would like to thank E. Dimastrogiovanni for drawing my attention to this error. In these versions the coefficient which accompanies $\ln k$ was also given incorrectly, owing to a calculational error which was present in v1–v2 of Ref. [58]. This error was corrected from v3 onwards of that reference, to which the reader should refer for details.

^{||} There is nothing to prevent powers of $\ln(-k\eta_*)$ accompanying q-loop corrections such as Eq. (20). These are genuine quantum corrections to the time evolution, and we should not expect the classical δN formula to account for their resummation. However, since these terms are suppressed by powers of H^2 we assume that their contribution is small.

of the same physics. It does not matter which time of evaluation we pick, as long as we have the ability to perform the relevant calculations consistently.

In this paper, we take the view that for analytic calculations it is better to evaluate the correlators of the fields soon after horizon crossing, where the logarithmic term $\text{Ci}(-2k\eta_*)$, and any higher-order logarithms, are negligible. There are two key reasons to adopt this approach:

- (i) The q-loop correlators given in Eqs. (19) and (20) were derived on the assumption that slow-roll is a good approximation between the time of horizon crossing and the time of observation of the $\delta\phi$. At any given order in the loop expansion one encounters a potentially large number of diagrams, but many of these diagrams will be subdominant when slow-roll applies. For example, as discussed in Ref. [58], the q-loop correction to the two-point function has a diagrammatic expansion of the form

$$\text{loop correction} \supseteq \text{---} \text{---} \text{---} \text{---} \text{---} + \text{---} \text{---} \text{---} \text{---} \text{---} + \dots$$

Both of these diagrams are one-loop and therefore of second order in cosmological perturbation theory, which is an expansion in powers of $\mathcal{P}_* \simeq 10^{-10}$. However, the second diagram is suppressed by an extra factor of order $\sim \epsilon$, since each three-point vertex carries a factor of $\dot{\phi}/H$ [94] whereas the four-point vertex carries no suppression [95]. On the other hand, large logarithms such as $\text{Ci}(-2k\eta_*)$ can compensate for the smallness of ϵ . Where large logarithms are present the first diagram is dominated by the terms of *subleading* order in slow-roll, since it is these terms which are accompanied by large logarithmic factors [51, 52]. When terms such as $\text{Ci}(-2k\eta_*)$ are large one must be wary that the second diagram (among other possible sources of subleading slow-roll terms) does not also begin to contribute. As the logarithms become increasingly large, further diagrams at two-loops and beyond may also become relevant. Therefore, in order to compute reliably with a slow-roll truncated q-loop such as Eq. (20) one should pick the point of evaluation in such a way that large logarithmic terms do not appear. This means that the initial hypersurface on which we base the δN expansion should be close to the horizon crossing hypersurface. The slow-roll expansion is then safe between these two hypersurfaces, and we can compute the q-loop to very good accuracy by assuming it is dominated by the leading slow-roll part of the first diagram. For the two-point function this difficulty is not severe, since for moderate values of $\ln(-k\eta_*)$ one can account for the contribution of the second diagram, together with the subleading part of the first. However, for higher-order correlation functions this may become a practical consideration.

- (ii) Eqs. (19) and (20) are computed from integrals over η whose integrands are rapidly oscillating when $|k\eta| \gg 1$. This kills any contribution from very early times, leaving the integral dominated by times where $|k\eta| \lesssim 1$. The pre-factors in Eqs. (19) and (20) are derived by approximating slowly varying quantities such as H and ϵ

at the time of horizon crossing. This will be a good approximation if the correlator is observed just a few e-folds after horizon crossing but may be on the verge of breaking down if $\ln |k\eta_*|$ is large, since in general one would then expect H and η to evolve between horizon crossing and the time of observation. One can account for this evolution using the δN expansion.

In the remainder of this paper only the “early” initial slice is considered, where the correlator is evaluated just a few e-folds after horizon crossing.

4. One-loop renormalization of the ζ power spectrum

4.1. The loop correction

We are now in a position to apply these general principles to the object of central interest, the power spectrum $\langle \zeta(\mathbf{k}_1)\zeta(\mathbf{k}_2) \rangle$. After collecting terms in Eqs. (14)–(17), the connected part of the power spectrum can be written

$$\langle \zeta(\mathbf{k}_1)\zeta(\mathbf{k}_2) \rangle = (2\pi)^3 \delta(\mathbf{k}_1 + \mathbf{k}_2) \left\{ \left(\frac{\partial N}{\partial \phi_*} \right)^2 P_*(k) + \int \frac{d^3 q}{(2\pi)^3} \Sigma(\mathbf{k}, \mathbf{q}) \right\} \quad (21)$$

where Σ is defined by

$$\Sigma \equiv \frac{\partial N}{\partial \phi_*} \frac{\partial^2 N}{\partial \phi_*^2} B_*(k, |\mathbf{k} - \mathbf{q}|, q) + \frac{\partial N}{\partial \phi_*} \frac{\partial^3 N}{\partial \phi_*^3} P_*(k) P_*(q) + \frac{1}{2} \left(\frac{\partial^2 N}{\partial \phi_*^2} \right)^2 P_*(|\mathbf{k} - \mathbf{q}|) P_*(q). \quad (22)$$

In this formula $\mathbf{k}$ can be taken to be either $\mathbf{k}_1$ or $\mathbf{k}_2$, since its orientation is immaterial, and the integration $d^3 q$ is restricted to momenta between $q \sim k$ and $q \sim \ell^{-1}$. As discussed above, the time η_* at which we construct the initial slice for the δN expansion is taken to be the horizon-crossing time for $\mathbf{k}$.

To estimate the magnitude of the loop correction in (21) we must be able to evaluate the $\mathbf{q}$ integral, but this is not trivial. In general, $B(k_1, k_2, k_3)$ is a complicated function of its arguments [94, 96] and the integral is difficult to evaluate exactly. We can obtain a reasonable estimate by applying the method of Boubekeur & Lyth [71]. In this approximation, one supposes that the integral is dominated by its infrared singularities. Summing over all singularities, one finds that

$$P_\zeta(k) = P_*(k) \left\{ \left(\frac{\partial N}{\partial \phi_*} \right)^2 + \Pi \right\}, \quad (23)$$

where P_* is the tree-level power spectrum evaluated at horizon crossing and Π is defined by

$$\Pi \equiv \left\{ \frac{\partial N}{\partial \phi_*} \frac{\partial^3 N}{\partial \phi_*^3} + \left(\frac{\partial^2 N}{\partial \phi_*^2} \right)^2 \right\} A(k\ell) - \frac{24}{5} f_{\text{NL}} \frac{\partial N}{\partial \phi_*} \frac{\partial^2 N}{\partial \phi_*^2} B(k\ell), \quad (24)$$

in which, as mentioned above, we have dropped the ultraviolet loop correction of Eq. (20). The functions A and B are defined by

$$A \equiv \langle \delta \phi^2 \rangle = \int \frac{d^3 q}{(2\pi)^3} P_*(q) \quad \text{and} \quad B \equiv \langle f_{\text{NL}} \delta \phi^2 \rangle = \int \frac{d^3 q}{(2\pi)^3} f_{\text{NL}} P_*(q). \quad (25)$$

The central issue in the remainder of this section, which will also be important for the background renormalization described in §4.2 below, is how one should obtain estimates of A and B .

4.2. The approximation of constant tilt

First assume that the power spectrum and f_{NL} can be approximated by Eq. (7) with constant tilt parameters $n - 1$ and n_{NL} . The A and B functions now satisfy

$$A(k\ell) = \mathcal{P}_* y_{n-1}(k\ell) \quad \text{and} \quad B(k\ell) = \mathcal{P}_* y_{n_{\text{NL}}+n-1}(k\ell). \quad (26)$$

where $y_r(x)$ is Lyth's y -function [72], defined by

$$y_r(x) \equiv \frac{1}{r} (1 - |x|^{-r}). \quad (27)$$

The parameters f_{NL} , n_{NL} and $n - 1$ are evaluated at k . Eq. (26) is approximately valid provided that the box size is not too much larger than the lengthscale associated with the mode of interest, k^{-1} , so that the approximation of fixed tilt is reasonable throughout the box.

The loop correction is of second order in cosmological perturbation theory, which is an expansion in powers of $\mathcal{P}$. However, the coefficients in Eq. (24) generated by the δN expansion enter at different orders in the slow-roll approximation. In particular, $(N')^2$ is of order ϵ^{-1} whereas $(N'')^2 + N'N'''$ is of order 1 and $f_{\text{NL}}N'N''$ is of order $\epsilon^{1/2}$. Which of these terms is dominant depends critically on the tilts $n - 1$ and n_{NL} .

- (i) *Positive tilt.* This is the case where the spectrum *decreases* on large scales, also known as a blue spectrum. It is not generically produced by single-field models of inflation. For positive r , the y -function defined in Eq. (27) satisfies $y_r(x) \sim 1/r$. If the spectral and bispectral tilts $n - 1$ and n_{NL} are both positive, then the loop correction is very small.
- (ii) *No tilt.* For $r = 0$, the y -function behaves like $y_r(x) \sim \ln|x|$. If $n - 1 = n_{\text{NL}} = 0$, it follows that all terms in Π behave like logarithms of $k\ell$. To estimate its magnitude, suppose that the loop correction constitutes no more than some fraction $f\%$ of the tree-level. It follows that

$$\Pi = \mathcal{P}_*(\Delta N - N_*) \approx \mathcal{P}_*\Delta N \lesssim \frac{f}{100}, \quad (28)$$

where ΔN is the total number of e-foldings between the scale ℓ and the end of inflation, and $N_* \approx 60$ measures when the presently observable universe left the horizon in terms of e-folds before inflation ended. The second approximate equality is valid if the loop correction is large, since this entails $\Delta N \gg N_*$. Eq. (28) can be interpreted as a bound on ΔN in order that the loop correction does not significantly modify the tree-level prediction. This gives $\Delta N \lesssim (10^2 f^{-1} \mathcal{P}_*)^{-1}$, or $\Delta N \lesssim 10^8$ for $f \sim 1\%$.

- (iii) *Negative tilt.* This is the case where the spectrum increases on large scales, also known as a red spectrum. It is the generic type of spectrum produced by single-field

models of inflation. For $r < 0$, the y -function has asymptotic form $y_r(x) \sim x^{|r|}/|r|$ when $x \gg 1$. In this regime, which term dominates depends whether $n_{\text{NL}} \geq 0$: the term involving f_{NL} will be dominant if n_{NL} is negative, meaning that the non-gaussianity grows on large scales.[†] For the purposes of making an estimate, however, it does not matter which term is the dominant one since in either case $\Pi \sim (k\ell)^t$, where $t > 0$ is the dominant tilt, and the difference in pre-factor is not significant. It follows that

$$\Pi \sim \frac{\alpha}{t} \mathcal{P}_* e^{Nt}, \quad (29)$$

where α is a numerical constant which absorbs the pre-factor. Eq. (29) gives an extremely powerful constraint on how strongly $\mathcal{P}$ and f_{NL} can tilt, but unfortunately it is untrustworthy precisely in the interesting limit where $k\ell \gg 1$ or if the tilt is significant. As discussed in §2, in this limit the approximation of constant tilt overpredicts the spectrum. In reality the spectrum does not grow like a constant power, but only a power of a logarithm.

Therefore we conclude that the significance of the loop correction depends on the number of e-folds of inflation between the fundamental scale, ℓ , and the scale of interest, k^{-1} . Except where there is a significant negative tilt the loop correction is small, perhaps of order 1% in a model with 10^7 e-folds of inflation. However, to study the case of negative tilt a more sophisticated analysis is required.

4.3. Monomial potentials.

To obtain a better prediction for $k\ell \gg 1$, or where there is negative tilt, consider any local field theory model of inflation, with potential given by the monomial [97]

$$V(\phi) = g\phi^r \quad (30)$$

where g is a coupling constant and r is usually taken to be an integer. For $r = 2$ and small values of the field this corresponds to an Nflation-type scenario [98], in which the quadratic behaviour arises as an approximation to a periodic axion potential. For larger r or large fields such monomials fit into the general framework of so-called chaotic inflation, where ϕ is taken to be descending from above the Planck scale with a spectrum of field values which may cause some regions of the universe to inflate at the expense of other regions in which the field value is not suitable for inflation to occur.

Inflation occurs for field values greater than $\phi_f \equiv r/\sqrt{2}$. However, one cannot contemplate arbitrarily large values for ϕ . Local field theory presumably breaks down as an approximate description of nature at least when $V \approx 1$ (in Planck units), which occurs at a field value $\phi_P \equiv g^{-1/r}$. This is certainly the largest value of the field for which the tools used in the present analysis make sense, but there may be other effects which imply that our approximations break down even before one reaches the Planck scale. For

[†] This is not the usual situation, since in single-field slow-roll inflation $f_{\text{NL}} \propto \epsilon$ [61], and therefore the non-gaussianity decays on large scales. At the times when the fluctuation on large scales was imprinted, ϵ was much closer to zero.

example, the ‘self-reproduction’ phase of eternal inflation ends when $2\delta\phi^2 \gtrsim 2\epsilon/3$ [97], which implies a maximum field value $\phi_{\text{sr}} \approx (2\pi^2 r^2 g^{-1})^{1/(r+2)}$. One can imagine either of these field values as natural energy scales at which to specify the masses and coupling constants of an effective field theory which approximates some ultraviolet completion of the Standard Model plus gravity.

Let us restrict attention to a regime where local field theory is a good approximation. Any two arbitrary field values ϕ_a and ϕ_b separated by ΔN e-folds of inflation are related via the rule

$$\phi_a^2 = \phi_b^2 + 2r\Delta N. \quad (31)$$

In order to match the fluctuation amplitude $\mathcal{P}_\zeta = (24.1 \pm 1.3) \times 10^{-10}^\ddagger$ observed in CMB experiments, the coupling must be tuned to precisely satisfy

$$g = \frac{12\pi^2 r^2 \mathcal{P}_\zeta}{\phi_*^{r+2}}, \quad (32)$$

where ϕ_* is the field value when presently observable scales were leaving the horizon. This typically occurs roughly $N_* \approx 60$ e-folds before the end of inflation in order that the flatness and horizon problems of classical cosmology are resolved. Eqs. (31)–(32) therefore imply a very small coupling constant, which is hard to understand at the Planck scale on the basis of our present knowledge of physics.

Suppose that inflation begins at some field value ϕ_{UV} . After a few e-foldings of inflation, we can suppose that the flat FRW metric (1) applies within the inflating volume. One therefore sets the box size ℓ to correspond to the horizon size at this time. The field value when any smaller scale q exits the horizon is related to the field value at the onset of inflation by (31), giving§

$$\phi^2(q) = \phi_{\text{UV}}^2 - 2r \ln q\ell. \quad (33)$$

It follows that $\langle \delta\phi^2 \rangle$ can be written [99, 51]

$$A \equiv \langle \delta\phi^2 \rangle = \frac{r}{r+4} \mathcal{P}_\zeta \left\{ \left(\frac{\phi_{\text{UV}}}{\phi_*} \right)^{r+4} - 1 \right\} \approx \frac{r}{r+4} \mathcal{P}_\zeta \left(\frac{\phi_{\text{UV}}}{\phi_*} \right)^{r+4}, \quad (34)$$

where the second approximate equality applies in the limit $\phi_* \ll \phi_{\text{UV}}$. If inflation lasts for a large total number of e-folds $N \gg r$, then Eqs. (31) and (34) imply

$$A = \frac{r}{r+4} \mathcal{P}_\zeta \left(\frac{N}{N_*} \right)^{2+r/2}. \quad (35)$$

This scaling of $\langle \delta\phi^2 \rangle \equiv A$ with the total number of e-folds, namely $A \propto N^{2+r/2}$, was given earlier by Sloth [52]. A similar procedure allows us to estimate B , giving

$$B = \frac{r^3}{r+2} \frac{\mathcal{P}_\zeta}{\phi_*^2} \left\{ \left(\frac{\phi_{\text{UV}}}{\phi_*} \right)^{r+2} - 1 \right\} \approx \frac{r^2}{2N_*(r+2)} \mathcal{P}_\zeta \left(\frac{N}{N_*} \right)^{1+r/2}. \quad (36)$$

[‡] The quoted value is for WMAP3+SDSS, ignoring any possible tensor contribution [20].

[§] In fact, ϕ_{UV} should be reduced by a small amount since we are assuming that ℓ does not correspond precisely to the inflationary patch when $\phi = \phi_{\text{UV}}$ but to some slightly later time. However, if the field is in slow-roll at this time then ϕ will not roll far during the few e-folds which elapse.

SIGNIFICANCE	$r = 2$ model			$r = 4$ model		
	maximum N	N_{P}	N_{sr}	maximum N	N_{P}	N_{sr}
0.1%	4.06×10^4			8.65×10^3		
1%	8.74×10^4	1.28×10^{10}	5.03×10^5	1.54×10^4	6.31×10^5	2.50×10^4
10%	1.88×10^5			2.74×10^4		

Table 1. Limits on the total number of e-folds, N , in monomial models with $r = 2$ and $r = 4$. N_{P} measures the number of e-folds between the Planck scale and the time presently observable scales were leaving the horizon, and N_{sr} is a similar measure, with the Planck scale replaced by the ‘self-reproduction’ scale of eternal inflation, ϕ_{sr} (defined in the text).

In making this estimate, we have approximated $f_{\text{NL}} \sim \epsilon$ at each scale. Although this is not strictly numerically accurate [61], it should be sufficient to capture the tilt of the non-gaussian fraction. Indeed, Eqs. (35)–(36) show that A grows faster than B by a power of N , and therefore that the non-gaussian component is negligible for a large total number of e-folds. This is in agreement with our expectation that the contribution to the loop correction from any intrinsic non-gaussianity among the $\delta\phi$ should be suppressed if f_{NL} decays on large scales. In addition, since both A and B only grow as powers of N , these equations justify the statements made above that a constant tilt overpredicts $\langle\delta\phi^2\rangle$ for a large range of e-folds. In what follows we assume that N is sufficiently large that it is a good approximation to ignore the intrinsically non-gaussian contribution.

Let us estimate how many e-folds can elapse before the loop correction becomes significant enough to spoil the tree-level prediction. The coefficient of the A -term in Eq. (24) is $\mathcal{O}(1)$ in slow-roll,

$$\frac{\partial N}{\partial\phi_*} \frac{\partial^3 N}{\partial\phi_*^3} + \left(\frac{\partial^2 N}{\partial\phi_*^2} \right)^2 = \frac{1}{r^2}. \quad (37)$$

Thus, in order that the loop contributes at no more than some fraction f of the tree-level, we must require $\Pi(\partial N/\partial\phi_*)^{-2} \lesssim f$, or

$$N \lesssim N_*^{(6+r)/(4+r)} \mathcal{P}_\zeta^{-2/(4+r)} \{2(4+r)f\}^{2/(4+r)}. \quad (38)$$

This gives constraints on N which are summarised in Table 1. Note that the bounds arising from the loop contribution are roughly compatible with the bounds derived by Wu *et al.* [26] on the basis that amplification of the quantum geometry should not spoil scale-invariance of the spectrum.

The numerical limits on N cannot be taken literally for **loop:tree** ratios as large as 10%, because the estimates for g and $\langle\delta\phi^2\rangle$ [given in Eqs. (32) and (34)] were derived assuming that the tree-level prediction dominated the power spectrum, and this is no longer true when the loop constitutes an appreciable fraction of the tree-level. Likewise, as the significance of the loop increases, contributions from two-loop

|| Note that in a monomial model there is a special simplification, since $N''' = 0$.

corrections and beyond may become relevant. Nevertheless, these limits show clearly that if inflation is taken to persist for an exponentially long time, corresponding to initial conditions at the Planck scale or the ‘self-reproduction’ scale of eternal inflation, then the loop correction will be large and cannot be ignored. On the other hand, if inflation is only taken to consist of sufficient e-folds to solve the horizon and flatness problems, $N \sim N_* \approx 60$, then the loop correction will be completely negligible. These conclusions are in qualitative agreement with the results of Sloth [52], who found corrections of $O(1)$ for inflation beginning at the self-reproduction scale. The qualitative disagreement between Refs. [51, 52] and the present paper on the one hand and the conclusions of Martineau & Brandenberger [87] on the other arises because in Ref. [87] the random distribution of phases present in the inflationary perturbations caused an overall cancellation of the effect. In the present formalism the distribution of phases is invisible; only the spectrum is relevant, not the modes themselves, and the large effect described in Table 1 is effectively the phase-space volume divergence given in Eq. (46) of Martineau & Brandenberger [87].

5. Discussion

In this paper, the δN formalism has been used to estimate the magnitude of the one-loop correction to the inflationary power spectrum in the case of a single-field, slow-roll model of the inflationary epoch. The calculation can be split into two separate parts. The first depends on a detailed estimate of the one-loop correction generated by interference among quantized field modes around the time of horizon crossing. This calculation has recently been performed in Ref. [58]. The second measures the pile-up of field modes outside the horizon which are generated by the quasi-de Sitter expansion. This accumulation of modes as inflation proceeds gives an ever-larger phase space for infrared divergences [87], which is measured by the so-called c-loops of the δN formalism. In this paper these two parts are assembled, allowing an estimate of the magnitude of the loop correction which includes all relevant effects.

The results is entirely consistent with our expectations from flat space quantum field theory: we are free to formulate the fundamental microscopic theory which governs inflation at any energy scale we wish, but if we choose to formulate it at an energy scale which is wildly different from that of the phenomena we wish to observe, then we must expect significant corrections to be generated by loop effects. Such corrections take the form of large logarithms that violate the predictions of naïve dimensional analysis. In the present case the loop correction is given by Eqs. (23)–(24), the fundamental theory is formulated at a scale ℓ , and we wish to predict phenomena associated with the CMB at a scale k^{-1} . The large logarithm which gives a significant loop correction is $N \sim \ln k\ell$. If $k\ell$ is exponentially large, which occurs in monomial models of inflation if initial conditions are set at around the Planck scale, then the loop correction has a sensitivity to the tilt of the field perturbation power spectrum. The δN loop from the convolution of two power spectra is the most sensitive and scales like $N^{2+n/2}$; the δN loop from any

intrinsic non-gaussianity among the $\delta\phi$ is suppressed, because f_{NL} scales in the opposite direction, giving $N^{1+n/2}$. In the case of a negative tilt, which produces perturbations of increasing amplitude on *large* scales, the loop correction rapidly becomes very significant. The correction is irrelevant only if inflation lasts for a small number of e-folds.

The physical effect which demands that loop corrections be taken into account is nothing more than back reaction. Long ago, Salopek & Bond [100] used the stochastic formalism to show that any initially homogeneous background field would develop wild fluctuations on ultra-large scales as inflation progressed. The large loop correction encountered in the limit of a large box is another manifestation of this effect. It measures the dispersion generated by comparing a correlator calculated in one region of the universe with the same correlator calculated in a different region, perhaps where the background fields have very different values. The correlator calculated in the large box is therefore not measureable in any experiment: it involves comparing the CMB temperature which would be seen in an ensemble of different universes, and no physical observer can make such a comparison. For this reason, the correlation is a “computable” in the sense of Witten [23], but is not an observable.

This result has no consequences for how we make or report cosmological observations, but has very significant consequences for how such observations are compared with theoretical models. In models where the fundamental theory is not taken to be valid within an exponentially large region, we are entitled to entirely disregard the question of loop corrections, since in this case they yield a negligible shift in tree-level values. This class of models includes the familiar single-field examples where inflation ends when the field reaches a certain point on its potential. In such cases we know the conditions which lead to the end of inflation and we always have the option to work within a small box.[†] On the other hand, where isocurvature fields are present, these fields will generally acquire large fluctuations within the inflating region. If these fields play a role in ending inflation, then we may lose the option to compute within a small box, whereas the correlator within a large box may be afflicted with large loop corrections. This problem is perhaps most severe precisely for those models to which we presently attach the greatest interest – those motivated from fundamental particle physics. For example, one popular class of such models arise from the so-called landscape of string vacua [101]. In such models, one imagines the unwanted six dimensions predicted by string theory to be wrapped up on a Calabi–Yau three-fold which may be topologically quite non-trivial, as measured by the existence of a large number of homological cycles. These cycles can be wrapped by fluxes of p -form fields, giving rise to regions of the three-fold which have different values for the cosmological constant Λ , together with the other constants of nature [102], and in general many scalar fields will be present which measure the size and shape of the Calabi–Yau three-

[†] Indeed, in such a case, it would presumably make no difference even if we chose to work within the large box because the only distinction between widely separated locations would be a time delay. This can be gauged away by a reparametrization of time, meaning that the difference is not physical. This is the essentially the argument given by Unruh [35].

fold. These different regions may exhibit substantially different low-energy physics. In such models, the natural scale at which one computes the masses and couplings in the low-energy theory would be the Planck scale, or perhaps the self-reproduction scale of eternal inflation. Eqs. (23)–(24) show that to compare any such theory to observation, one must account for large loop corrections which are inevitably encountered in passing from the Planck scale to the scale relevant for CMB predictions.

This has some implications for how large-field models of inflation are compared with CMB observations. To understand how the Planck scale theory is related to the theory when scales associated with the CMB exited the horizon, one would need to integrate out the relevant degrees of freedom between M_P and the CMB scale. By analogy with conventional field theory, as one changes scale the effective lagrangian should be controlled by a flow equation analogous to Polchinski’s exact renormalization group equation [103]. In flat space theory this flow has well-understood effects, attracting the lagrangian to an invariant hypersurface on which non-renormalizable couplings are suppressed. (This would refer to the effective field theory for ζ , not to the couplings in the matter theory.) Unfortunately a complete understanding of this process in the cosmological case is not yet in hand, and it does not appear to be understood precisely how the Planck scale theory will be related to the theory in a small box. One possible approach is to use the method of Salopek & Bond [100], who gave a Fokker–Planck equation which determined the probability distribution of the classical fields on large scales. Having done so, it is possible to use a small box to determine the value of the correlation functions at each value of the allowed background fields, but one is then restricted to making probabilistic statements concerning what should be observed in the CMB. This issue deserves further study.

In any case, considerable work remains to be done. The present analysis should be generalized to the case of multiple scalar fields in order that it can be applied to the best-motivated models from fundamental physics. This involves an extension of the q-loop calculation given in Ref. [58] to the case of many scalar fields. The calculation given by Weinberg [44] shows that where ζ is accompanied by $\mathcal{N}$ isocurvature fields, we can expect the ζ loop to receive an enhancement by a factor $\sim \mathcal{N}$. However, in a multiple field scenario a more involved process is required to estimate $\langle \delta\phi^2 \rangle$ [104, 105, 106, 107], since the growth of $\mathcal{P}$ on superhorizon scales is no longer simple to predict. These issues should be addressed with some urgency.

Acknowledgments

I acknowledge support from PPARC under grant PPA/G/S/2003/00076. I would like to thank C. Byrnes, J. Lidsey, D. Lyth, K. Malik, A. Mazumdar, A. Riotto and M. Sloth for useful conversations, and especially F. Vernizzi for lengthy conversations and correspondence which have helped clarify my understanding. I would like to acknowledge the hospitality of the Abdus Salam Institute for Theoretical Physics, Trieste, and the Department of Physics, University of Cardiff, where portions of the work outlined in

this paper were carried out. I would like to thank the Yukawa Institute for Theoretical Physics, Kyoto, for their hospitality while participating in the programme *Gravity and Cosmology 2007*, where this paper was completed.

I would like to thank E. Dimastrogiovanni for drawing my attention to an error present in earlier versions of this paper (discussed below Eq. (20)).

References

- [1] A. A. Starobinsky, *A new type of isotropic cosmological models without singularity*, *Phys. Lett.* **B91** (1980) 99–102.
- [2] K. Sato, *First order phase transition of a vacuum and expansion of the universe*, *Mon. Not. Roy. Astron. Soc.* **195** (1981) 467–479.
- [3] A. H. Guth, *The inflationary universe: A possible solution to the horizon and flatness problems*, *Phys. Rev.* **D23** (1981) 347–356.
- [4] S. W. Hawking and I. G. Moss, *Supercooled phase transitions in the very early universe*, *Phys. Lett.* **B110** (1982) 35.
- [5] A. Albrecht and P. J. Steinhardt, *Cosmology for grand unified theories with radiatively induced symmetry breaking*, *Phys. Rev. Lett.* **48** (1982) 1220–1223.
- [6] A. D. Linde, *A new inflationary universe scenario: A possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems*, *Phys. Lett.* **B108** (1982) 389–393.
- [7] A. D. Linde, *Chaotic inflation*, *Phys. Lett.* **B129** (1983) 177–181.
- [8] J. M. Bardeen, P. J. Steinhardt, and M. S. Turner, *Spontaneous creation of almost scale-free density perturbations in an inflationary universe*, *Phys. Rev.* **D28** (1983) 679.
- [9] A. H. Guth and S. Y. Pi, *Fluctuations in the new inflationary universe*, *Phys. Rev. Lett.* **49** (1982) 1110–1113.
- [10] S. W. Hawking, *The development of irregularities in a single bubble inflationary universe*, *Phys. Lett.* **B115** (1982) 295.
- [11] S. W. Hawking and I. G. Moss, *Fluctuations in the inflationary universe*, *Nucl. Phys.* **B224** (1983) 180.
- [12] V. F. Mukhanov, *Gravitational instability of the universe filled with a scalar field*, *JETP Lett.* **41** (1985) 493–496.
- [13] M. Sasaki, *Large scale quantum fluctuations in the inflationary universe*, *Prog. Theor. Phys.* **76** (1986) 1036.
- [14] V. F. Mukhanov, *Quantum theory of cosmological perturbations in R^2 gravity*, *Phys. Lett.* **B218** (1989) 17–20.
- [15] V. F. Mukhanov, H. A. Feldman, and R. H. Brandenberger, *Theory of cosmological perturbations. Part 1. Classical perturbations. Part 2. Quantum theory of perturbations. Part 3. Extensions*, *Phys. Rept.* **215** (1992) 203–333.
- [16] D. H. Lyth and D. Seery, *Classicality of the primordial perturbations*, [arXiv:astro-ph/0607647](#).
- [17] D. H. Lyth and D. Wands, *Generating the curvature perturbation without an inflaton*, *Phys. Lett.* **B524** (2002) 5–14, [[arXiv:hep-ph/0110002](#)].
- [18] T. Moroi and T. Takahashi, *Effects of cosmological moduli fields on cosmic microwave background*, *Phys. Lett.* **B522** (2001) 215–221, [[arXiv:hep-ph/0110096](#)].
- [19] S. Mollerach, *Isocurvature baryon perturbations and inflation*, *Phys. Rev.* **D42** (1990) 313–325.
- [20] **WMAP** Collaboration, D. N. Spergel *et al.*, *Wilkinson Microwave Anisotropy probe (WMAP) three year results: Implications for cosmology*, [arXiv:astro-ph/0603449](#).
- [21] J. Martin and C. Ringeval, *Inflation after WMAP3: Confronting the slow-roll and exact power spectra to CMB data*, *JCAP* **0608** (2006) 009, [[arXiv:astro-ph/0605367](#)].
- [22] W. H. Kinney, E. W. Kolb, A. Melchiorri, and A. Riotto, *Inflation model constraints from the*

- Wilkinson Microwave Anisotropy Probe three-year data*, *Phys. Rev.* **D74** (2006) 023502, [arXiv:astro-ph/0605338].
- [23] E. Witten, *Quantum gravity in de Sitter space*, arXiv:hep-th/0106109.
 - [24] N. Arkani-Hamed, S. Dubovsky, A. Nicolis, E. Trincherini, and G. Villadoro, *A measure of de Sitter entropy and eternal inflation*, *JHEP* **05** (2007) 055, [arXiv:arXiv:0704.1814 [hep-th]].
 - [25] S. B. Giddings and D. Marolf, *A global picture of quantum de Sitter space*, arXiv:arXiv:0705.1178 [hep-th].
 - [26] C.-H. Wu, K.-W. Ng, and L. H. Ford, *Possible constraints on the duration of inflationary expansion from quantum stress tensor fluctuations*, *Phys. Rev.* **D75** (2007) 103502, [arXiv:gr-qc/0608002].
 - [27] B. Losic and W. G. Unruh, *On leading order gravitational backreactions in de Sitter spacetime*, *Phys. Rev.* **D74** (2006) 023511, [arXiv:gr-qc/0604122].
 - [28] Q.-G. Huang, M. Li, and Y. Wang, *Eternal chaotic inflation is prohibited by weak gravity conjecture*, arXiv:arXiv:0707.3471 [hep-th].
 - [29] G. 't Hooft and M. J. G. Veltman, *One loop divergencies in the theory of gravitation*, *Annales Poincare Phys. Theor.* **A20** (1974) 69–94.
 - [30] M. Sasaki, H. Suzuki, K. Yamamoto, and J. Yokoyama, *Superexpansionary divergence: Breakdown of perturbative quantum field theory in space-time with accelerated expansion*, *Class. Quant. Grav.* **10** (1993) L55–L60.
 - [31] H. Suzuki, M. Sasaki, K. Yamamoto, and J. Yokoyama, *Probability distribution functional for equal time correlation functions in curved space*, *Int. J. Mod. Phys.* **A9** (1994) 221–238.
 - [32] V. F. Mukhanov, L. R. W. Abramo, and R. H. Brandenberger, *On the back reaction problem for gravitational perturbations*, *Phys. Rev. Lett.* **78** (1997) 1624–1627, [arXiv:gr-qc/9609026].
 - [33] L. R. W. Abramo, R. H. Brandenberger, and V. F. Mukhanov, *The energy-momentum tensor for cosmological perturbations*, *Phys. Rev.* **D56** (1997) 3248–3257, [arXiv:gr-qc/9704037].
 - [34] L. R. W. Abramo and R. P. Woodard, *One loop back reaction on chaotic inflation*, *Phys. Rev.* **D60** (1999) 044010, [arXiv:astro-ph/9811430].
 - [35] W. Unruh, *Cosmological long wavelength perturbations*, arXiv:astro-ph/9802323.
 - [36] T. Prokopec, O. Tornkvist, and R. P. Woodard, *One loop vacuum polarization in a locally de Sitter background*, *Ann. Phys.* **303** (2003) 251–274, [arXiv:gr-qc/0205130].
 - [37] V. K. Onemli and R. P. Woodard, *Super-acceleration from massless, minimally coupled ϕ^4* , *Class. Quant. Grav.* **19** (2002) 4607, [arXiv:gr-qc/0204065].
 - [38] T. Prokopec and R. P. Woodard, *Vacuum polarization and photon mass in inflation*, *Am. J. Phys.* **72** (2004) 60–72, [arXiv:astro-ph/0303358].
 - [39] V. K. Onemli and R. P. Woodard, *Quantum effects can render $w < -1$ on cosmological scales*, *Phys. Rev.* **D70** (2004) 107301, [arXiv:gr-qc/0406098].
 - [40] T. Brunier, V. K. Onemli, and R. P. Woodard, *Two loop scalar self-mass during inflation*, *Class. Quant. Grav.* **22** (2005) 59–84, [arXiv:gr-qc/0408080].
 - [41] D. Boyanovsky, H. J. de Vega, and N. G. Sanchez, *Quantum corrections to slow roll inflation and new scaling of superhorizon fluctuations*, *Nucl. Phys.* **B747** (2006) 25–54, [arXiv:astro-ph/0503669].
 - [42] D. Boyanovsky, H. J. de Vega, and N. G. Sanchez, *Quantum corrections to the inflaton potential and the power spectra from superhorizon modes and trace anomalies*, *Phys. Rev.* **D72** (2005) 103006, [arXiv:astro-ph/0507596].
 - [43] E. O. Kahya and R. P. Woodard, *Charged scalar self-mass during inflation*, *Phys. Rev.* **D72** (2005) 104001, [arXiv:gr-qc/0508015].
 - [44] S. Weinberg, *Quantum contributions to cosmological correlations*, *Phys. Rev.* **D72** (2005) 043514, [arXiv:hep-th/0506236].
 - [45] K. Chaicherdsakul, *Quantum cosmological correlations in an inflating universe: Can fermion and gauge fields loops give a scale free spectrum?*, *Phys. Rev.* **D75** (2007) 063522, [arXiv:hep-th/0611352].

- [46] B. Losic and W. G. Unruh, *Long-wavelength metric backreactions in slow-roll inflation*, *Phys. Rev.* **D72** (2005) 123510, [arXiv:gr-qc/0510078].
- [47] E. O. Kahya and V. K. Onemli, *Quantum stability of a $w < -1$ phase of cosmic acceleration*, arXiv:gr-qc/0612026.
- [48] A. Bilandžić and T. Prokopec, *Quantum radiative corrections to slow-roll inflation*, arXiv:arXiv:0704.1905 [astro-ph].
- [49] E. O. Kahya and R. P. Woodard, *One loop corrected mode functions for SQED during inflation*, *Phys. Rev.* **D74** (2006) 084012, [arXiv:gr-qc/0608049].
- [50] T. Prokopec, N. C. Tsamis, and R. P. Woodard, *Two loop scalar bilinears for inflationary SQED*, *Class. Quant. Grav.* **24** (2007) 201–230, [arXiv:gr-qc/0607094].
- [51] M. S. Sloth, *On the one loop corrections to inflation and the CMB anisotropies*, *Nucl. Phys.* **B748** (2006) 149–169, [arXiv:astro-ph/0604488].
- [52] M. S. Sloth, *On the one loop corrections to inflation. II: The consistency relation*, *Nucl. Phys.* **B775** (2007) 78–94, [arXiv:hep-th/0612138].
- [53] S. P. Kim, *Quantum fluctuations in the inflationary universe*, arXiv:astro-ph/0701399.
- [54] A. A. Starobinsky, *Multicomponent de Sitter (inflationary) stages and the generation of perturbations*, *JETP Lett.* **42** (1985) 152–155.
- [55] M. Sasaki and E. D. Stewart, *A general analytic formula for the spectral index of the density perturbations produced during inflation*, *Prog. Theor. Phys.* **95** (1996) 71–78, [arXiv:astro-ph/9507001].
- [56] D. H. Lyth, K. A. Malik, and M. Sasaki, *A general proof of the conservation of the curvature perturbation*, *JCAP* **0505** (2005) 004, [arXiv:astro-ph/0411220].
- [57] D. H. Lyth and Y. Rodríguez, *The inflationary prediction for primordial non-gaussianity*, *Phys. Rev. Lett.* **95** (2005) 121302, [arXiv:astro-ph/0504045].
- [58] D. Seery, *One-loop corrections to a scalar field during inflation*, arXiv:arXiv:0707.3377 [astro-ph].
- [59] N. Bartolo, E. Komatsu, S. Matarrese, and A. Riotto, *Non-gaussianity from inflation: Theory and observations*, *Phys. Rept.* **402** (2004) 103–266, [arXiv:astro-ph/0406398].
- [60] E. Komatsu and D. N. Spergel, *Acoustic signatures in the primary microwave background bispectrum*, *Phys. Rev.* **D63** (2001) 063002, [arXiv:astro-ph/0005036].
- [61] J. M. Maldacena, *Non-gaussian features of primordial fluctuations in single field inflationary models*, *JHEP* **05** (2003) 013, [arXiv:astro-ph/0210603].
- [62] X. Chen, *Running non-gaussianities in DBI inflation*, *Phys. Rev.* **D72** (2005) 123518, [arXiv:astro-ph/0507053].
- [63] A. Kosowsky and M. S. Turner, *CMB anisotropy and the running of the scalar spectral index*, *Phys. Rev.* **D52** (1995) 1739–1743, [arXiv:astro-ph/9504071].
- [64] S. Weinberg, *Quantum contributions to cosmological correlations. II: Can these corrections become large?*, *Phys. Rev.* **D74** (2006) 023508, [arXiv:hep-th/0605244].
- [65] J. M. Bardeen, *Gauge invariant cosmological perturbations*, *Phys. Rev.* **D22** (1980) 1882–1905.
- [66] G. I. Rigopoulos and E. P. S. Shellard, *Non-linear inflationary perturbations*, *JCAP* **0510** (2005) 006, [arXiv:astro-ph/0405185].
- [67] R. Arnowitt, S. Deser, and C. W. Misner, *Canonical variables for general relativity*, *Phys. Rev.* **117** (1960) 1595–1602.
- [68] R. W. Wald, *Asymptotic behavior of homogeneous cosmological models in the presence of a positive cosmological constant*, *Phys. Rev.* **D28** (1983) 2118–2120.
- [69] J. C. Collins, *Renormalization – an introduction to renormalization, the renormalization group, and the operator product expansion*. Cambridge University Press, Cambridge, 1984. 380 p.
- [70] D. H. Lyth, *Axions and inflation: Sitting in the vacuum*, *Phys. Rev.* **D45** (1992) 3394–3404.
- [71] L. Boubekur and D. H. Lyth, *Detecting a small perturbation through its non-gaussianity*, *Phys. Rev.* **D73** (2006) 021301, [arXiv:astro-ph/0504046].
- [72] D. H. Lyth, *Non-gaussianity and cosmic uncertainty in curvaton-type models*, *JCAP* **0606** (2006) 015, [arXiv:astro-ph/0602285].

- [73] D. H. Lyth, *The curvature perturbation in a box*, arXiv:arXiv:0707.0361 [astro-ph].
- [74] M. Crocce and R. Scoccimarro, *Renormalized cosmological perturbation theory*, *Phys. Rev.* **D73** (2006) 063519, [arXiv:astro-ph/0509418].
- [75] I. Zaballa, Y. Rodríguez, and D. H. Lyth, *Higher order contributions to the primordial non-gaussianity*, *JCAP* **0606** (2006) 013, [arXiv:astro-ph/0603534].
- [76] C. T. Byrnes, K. Koyama, M. Sasaki, and D. Wands, *Diagrammatic approach to non-gaussianity from inflation*, arXiv:arXiv:0705.4096 [hep-th].
- [77] E. W. Kolb, S. Matarrese, and A. Riotto, *On cosmic acceleration without dark energy*, *New J. Phys.* **8** (2006) 322, [arXiv:astro-ph/0506534].
- [78] E. W. Kolb, S. Matarrese, A. Notari, and A. Riotto, *Primordial inflation explains why the universe is accelerating today*, arXiv:hep-th/0503117.
- [79] G. Geshnizjani, D. J. H. Chung, and N. Afshordi, *Do large-scale inhomogeneities explain away dark energy?*, *Phys. Rev.* **D72** (2005) 023517, [arXiv:astro-ph/0503553].
- [80] E. E. Flanagan, *Can superhorizon perturbations drive the acceleration of the universe?*, *Phys. Rev.* **D71** (2005) 103521, [arXiv:hep-th/0503202].
- [81] C. M. Hirata and U. Seljak, *Can superhorizon cosmological perturbations explain the acceleration of the universe?*, *Phys. Rev.* **D72** (2005) 083501, [arXiv:astro-ph/0503582].
- [82] A. Notari, *Late time failure of Friedmann equation*, *Mod. Phys. Lett.* **A21** (2006) 2997–3001, [arXiv:astro-ph/0503715].
- [83] S. Rasanen, *Backreaction and spatial curvature in a dust universe*, *Class. Quant. Grav.* **23** (2006) 1823–1835, [arXiv:astro-ph/0504005].
- [84] J. W. Moffat, *Large scale cosmological inhomogeneities, inflation and acceleration without dark energy*, arXiv:astro-ph/0504004.
- [85] R. H. Brandenberger, *Back reaction of cosmological perturbations and the cosmological constant problem*, arXiv:hep-th/0210165.
- [86] R. Brandenberger and A. Mazumdar, *Dynamical relaxation of the cosmological constant and matter creation in the universe*, *JCAP* **0408** (2004) 015, [arXiv:hep-th/0402205].
- [87] P. Martineau and R. H. Brandenberger, *The effects of gravitational back-reaction on cosmological perturbations*, *Phys. Rev.* **D72** (2005) 023507, [arXiv:astro-ph/0505236].
- [88] P. Martineau and R. Brandenberger, *Back-reaction: A cosmological panacea*, arXiv:astro-ph/0510523.
- [89] J.-O. Gong and E. D. Stewart, *The density perturbation power spectrum to second-order corrections in the slow-roll expansion*, *Phys. Lett.* **B510** (2001) 1–9, [arXiv:astro-ph/0101225].
- [90] J.-O. Gong and E. D. Stewart, *The power spectrum for a multi-component inflaton to second-order corrections in the slow-roll expansion*, *Phys. Lett.* **B538** (2002) 213–222, [arXiv:astro-ph/0202098].
- [91] A. A. Starobinsky and J. Yokoyama, *Equilibrium state of a selfinteracting scalar field in the de Sitter background*, *Phys. Rev.* **D50** (1994) 6357–6368, [arXiv:astro-ph/9407016].
- [92] N. C. Tsamis and R. P. Woodard, *Stochastic quantum gravitational inflation*, *Nucl. Phys.* **B724** (2005) 295–328, [arXiv:gr-qc/0505115].
- [93] S.-P. Miao and R. P. Woodard, *Leading log solution for inflationary Yukawa*, *Phys. Rev.* **D74** (2006) 044019, [arXiv:gr-qc/0602110].
- [94] D. Seery and J. E. Lidsey, *Primordial non-gaussianities from multiple-field inflation*, *JCAP* **0509** (2005) 011, [arXiv:astro-ph/0506056].
- [95] D. Seery, J. E. Lidsey, and M. S. Sloth, *The inflationary trispectrum*, *JCAP* **0701** (2007) 027, [arXiv:astro-ph/0610210].
- [96] L. E. Allen, S. Gupta, and D. Wands, *Non-gaussian perturbations from multi-field inflation*, *JCAP* **0601** (2006) 006, [arXiv:astro-ph/0509719].
- [97] A. D. Linde, D. A. Linde, and A. Mezhlumian, *From the big bang theory to the theory of a stationary universe*, *Phys. Rev.* **D49** (1994) 1783–1826, [arXiv:gr-qc/9306035].
- [98] S. Dimopoulos, S. Kachru, J. McGreevy, and J. G. Wacker, *N-flation*, arXiv:hep-th/0507205.

- [99] N. Afshordi and R. H. Brandenberger, *Super-Hubble nonlinear perturbations during inflation*, *Phys. Rev.* **D63** (2001) 123505, [arXiv:gr-qc/0011075].
- [100] D. S. Salopek and J. R. Bond, *Stochastic inflation and nonlinear gravity*, *Phys. Rev.* **D43** (1991) 1005–1031.
- [101] S. B. Giddings, S. Kachru, and J. Polchinski, *Hierarchies from fluxes in string compactifications*, *Phys. Rev.* **D66** (2002) 106006, [arXiv:hep-th/0105097].
- [102] R. Bousso and J. Polchinski, *Quantization of four-form fluxes and dynamical neutralization of the cosmological constant*, *JHEP* **06** (2000) 006, [arXiv:hep-th/0004134].
- [103] J. Polchinski, *Renormalization and effective Lagrangians*, *Nucl. Phys.* **B231** (1984) 269–295.
- [104] J. Garcia-Bellido and D. Wands, *Metric perturbations in two-field inflation*, *Phys. Rev.* **D53** (1996) 5437–5445, [arXiv:astro-ph/9511029].
- [105] F. Vernizzi and D. Wands, *Non-gaussianities in two-field inflation*, *JCAP* **0605** (2006) 019, [arXiv:astro-ph/0603799].
- [106] T. Battefeld and R. Easther, *Non-gaussianities in multi-field inflation*, *JCAP* **0703** (2007) 020, [arXiv:astro-ph/0610296].
- [107] S. Yokoyama, T. Suyama, and T. Tanaka, *Primordial non-gaussianity in multi-scalar slow-roll inflation*, arXiv:arXiv:0705.3178 [astro-ph].